
Product Information

Size:

50ug

Protein Background:

Minor protein of the capsid that localizes along the inner surface of the virion, within the central cavities beneath the L1 pentamers. Plays a role in capsid stabilization through interaction with the major capsid protein L1. Once the virion enters the host cell, escorts the genomic DNA into the nucleus, in particular by promoting virion endosomal escape. It is involved, through its interaction with host dynein, in the

Reactivity:

Human papillomavirus type 16

cell, escorts the genomic DNA into the nucleus, in particular by promoting virion endosomal escape. It is involved, through its interaction with host dynein, in the intracellular microtubule-dependent transport of viral capsid toward the nucleus.

Source:

Rabbit

Mediates the viral genome import into the nucleus through binding to host importins. Once within the nucleus, L2 localizes viral genomes to PML bodies in order to activate

Isotype:

IgG

early gene expression for establishment of infection. Later on, promotes late gene expression by interacting with the viral E2 protein and by inhibiting its transcriptional activation functions. During virion assembly, encapsidates the genome by direct interaction with the viral DNA.

Applications:

ELISA

Gene ID:

L2

Recommended dilutions:**Uniprot**

P03107

Synonyms:

Minor capsid protein L2, L2

Immunogen:

Recombinant Human papillomavirus type 16 Minor capsid protein L2 protein (1-473AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product Images

N/A

N/A