METRNL Antibody

PACO36574




Size:	Protein Background:
50ug	Hormone induced following exercise or cold exposure that promotes energy
Reactivity:	expenditure. Induced either in the skeletal muscle after exercise or in adipose tissue following cold exposure and is present in the circulation. Able to stimulate energy
Human, Mouse, Rat	expenditure associated with the browning of the white fat depots and improves glucose tolerance. Does not promote an increase in a thermogenic gene program via
Source:	direct action on adipocytes, but acts by stimulating several immune cell subtypes to
Rabbit	enter the adipose tissue and activate their prothermogenic actions. Stimulates an eosinophil-dependent increase in IL4 expression and promotes alternative activation of
lsotype:	adipose tissue macrophages, which are required for the increased expression of the thermogenic and anti-inflammatory gene programs in fat. Required for some cold-
lgG	induced thermogenic responses, suggesting a role in metabolic adaptations to cold
Applications:	temperatures.
ELISA, WB, IHC	Gene ID:
Recommended dilutions:	METRNL
	Uniprot
ELISA:1:2000-1:10000, WB:1:500-1:5000, IHC:1:20-1:200	Q641Q3
	Synonyms:
	Meteorin-like protein (Subfatin), METRNL
	Immunogen:
	Recombinant Human Meteorin-like protein (46-311AA).

Storage:


Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4





Western blot. All lanes: METRNL antibody at 2µg/ml. Lane 1: Mouse small intestine tissue. Lane 2: Mouse kidney tissue. Lane 3: Rat gonadal tissue. Lane 4: Mouse brain tissue. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 35, 26 kDa. Observed band size: 35 kDa.

Lane1 Lane2 Lane3 Lane4



Immunohistochemistry of paraffin-embedded human kidney tissue using PACO36574 at dilution of 1:100.