PEG10 Antibody, Biotin conjugated



PACO37229

Human

Source:

Product Information

Size: **Protein Background:**

50ug Prevents apoptosis in hepatocellular carcinoma (HCC) cells through interaction with SIAH1, a mediator of apoptosis. May also have a role in cell growth promotion and Reactivity: hepatoma formation. Inhibits the TGF-beta signaling by interacting with the TGF-beta

> receptor ALK1. When overexpressed, induces the formation of cellular extension, such as filipodia in association with ALK1. Involved at the immediate early stage of adipocyte

differentiation By similarity. May bind to the 5'-GCCTGTCTTT-3' DNA sequence of the MB1 domain in the myelin basic protein (MBP) promoter.

Rabbit

PEG10

Gene ID: Isotype:

lgG Uniprot

Applications: Q86TG7 **ELISA**

Synonyms: **Recommended dilutions:**

> Retrotransposon-derived protein PEG10 (Embryonal carcinoma differentiationregulated protein) (Mammalian retrotransposon-derived protein 2) (Myelin expression factor 3-like protein 1) (MEF3-like protein 1) (Paternally expressed gene 10 protein) (Retrotransposon gag domain-containing protein 3) (Retrotransposon-derived gag-like polyprotein) (Ty3/Gypsy-like protein), PEG10, EDR KIAA1051 MAR2 MART2 MEF3L1 RGAG3

Immunogen:

Recombinant Human Retrotransposon-derived protein PEG10 protein (1-325AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product	Images
---------	---------------

N/A N/A