

Product Information

Size:

50ug

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:1000-1:5000

Protein Background:

The production of the second messenger molecules diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. In vitro, hydrolyzes PtdIns(4,5)P2 in a Ca(2+)-dependent manner. Triggers intracellular Ca(2+) oscillations in oocytes solely during M phase and is involved in inducing oocyte activation and initiating embryonic development up to the blastocyst stage. Is therefore a strong candidate for the egg-activating soluble sperm factor that is transferred from the sperm into the egg cytoplasm following gamete membrane fusion. May exert an inhibitory effect on phospholipase-C-coupled processes that depend on calcium ions and protein kinase C, including CFTR trafficking and function.

Gene ID:

PLCZ1

Uniprot

Q86YW0

Synonyms:

1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase zeta-1 (EC 3.1.4.11) (Phosphoinositide phospholipase C-zeta-1) (Phospholipase C-zeta-1) (PLC-zeta-1) (Testis-development protein NYD-SP27), PLCZ1


Immunogen:

Recombinant Human 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase zeta-1 protein (1-415AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product Images



Western blot. All lanes: PLCZ1 antibody at 3 μ g/ml. Lane 1: Mouse liver tissue. Lane 2: Mouse kidney tissue. Lane 3: Human placenta tissue. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 71, 48, 58 kDa. Observed band size: 71 kDa.