

### Product Information

**Size:**

50ug

**Reactivity:**

Human

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, IHC, IF

**Recommended dilutions:**

ELISA:1:2000-1:10000, IHC:1:20-1:200,  
IF:1:50-1:500

**Protein Background:**

ATP-dependent RNA helicase. Binds to partially double-stranded RNAs (dsRNAs) in order to unwind RNA secondary structures. Unwinding is promoted in the presence of single-strand binding proteins. Mediates also RNA duplex formation thereby displacing the single-strand RNA binding protein. ATP and ADP modulate its activity: ATP binding and hydrolysis by DDX42 triggers RNA strand separation, whereas the ADP-bound form of the protein triggers annealing of complementary RNA strands. Involved in the survival of cells by interacting with TP53BP2 and thereby counteracting the apoptosis-stimulating activity of TP53BP2. Relocalizes TP53BP2 to the cytoplasm.

**Gene ID:**

DDX42

**Uniprot**

Q86XP3

**Synonyms:**

ATP-dependent RNA helicase DDX42 (EC 3.6.4.13) (DEAD box protein 42) (RNA helicase-like protein) (RHELP) (RNA helicase-related protein) (RNAHP) (SF3b DEAD box protein) (Splicing factor 3B-associated 125 kDa protein) (SF3b125), DDX42

**Immunogen:**


Recombinant Human ATP-dependent RNA helicase DDX42 protein (519-819AA).

**Storage:**


Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

## Product Images


---



Immunohistochemistry of paraffin-embedded human tonsil tissue using PACO37422 at dilution of 1:100.



Immunofluorescence staining of HeLa cells with PACO37422 at 1:200, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemistry of paraffin-embedded human lung tissue using PACO37422 at dilution of 1:100.