PTPMT1 Antibody, Biotin conjugated

PACO38057



Product Information	
Size:	Protein Background:
50ug	Lipid phosphatase which dephosphorylates phosphatidylglycerophosphate (PGP) to
Reactivity:	phosphatidylglycerol (PG). PGP is an essential intermediate in the biosynthetic pathway of cardiolipin, a mitochondrial-specific phospholipid regulating the membrane integrity
Human	and activities of the organelle. Has also been shown to display phosphatase activity toward phosphoprotein substrates, specifically mediates dephosphorylation of
Source:	mitochondrial proteins, thereby playing an essential role in ATP production. Has
Rabbit	probably a preference for proteins phosphorylated on Ser and/or Thr residues compared to proteins phosphorylated on Tyr residues. Probably involved in regulation
lsotype:	of insulin secretion in pancreatic beta cells.
lgG	Gene ID:
Applications:	PTPMT1
ELISA	Uniprot
Recommended dilutions:	Q8WUK0
	Synonyms:
	Phosphatidylglycerophosphatase and protein-tyrosine phosphatase 1 (EC 3.1.3.27) (PTEN-like phosphatase) (Phosphoinositide lipid phosphatase) (Protein-tyrosine phosphatase mitochondrial 1) (EC 3.1.3.16) (EC 3.1.3.48), PTPMT1, MOSP PLIP
	Immunogen:
	Recombinant Human Phosphatidylglycerophosphatase and protein-tyrosine phosphatase 1 protein (28-201AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A