HSD17B11 Antibody, HRP conjugated



PACO39127

Rabbit

Applications:

Product Information

Size: **Protein Background:**

50ug Can convert androstan-3-alpha,17-beta-diol (3-alpha-diol) to androsterone in vitro, suggesting that it may participate in androgen metabolism during steroidogenesis.

Reactivity: May act by metabolizing compounds that stimulate steroid synthesis and/or by

generating metabolites that inhibit it. Has no activity toward DHEA Human

(dehydroepiandrosterone), or A-dione (4-androste-3,17-dione), and only a slight Source:

activity toward testosterone to A-dione. Tumor-associated antigen in cutaneous T-cell

lymphoma.

Gene ID: Isotype:

HSD17B11

lgG Uniprot

Q8NBQ5 **ELISA**

Synonyms: **Recommended dilutions:**

> Estradiol 17-beta-dehydrogenase 11 (EC 1.1.1.62) (17-beta-hydroxysteroid dehydrogenase 11) (17-beta-HSD 11) (17bHSD11) (17betaHSD11) (17-betahydroxysteroid dehydrogenase XI) (17-beta-HSD XI) (17betaHSDXI) (Cutaneous T-cell lymphoma-associated antigen HD-CL-03) (CTCL-associated antigen HD-CL-03) (Dehydrogenase/reductase SDR family member 8) (Retinal short-chain dehydrogenase/reductase 2) (retSDR2) (Short chain dehydrogenase/reductase family 16C member 2), HSD17B11, DHRS8 PAN1B SDR16C2

Immunogen:

Recombinant Human Estradiol 17-beta-dehydrogenase 11 protein (20-300AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product	Images
---------	---------------

N/A N/A