

# PPIL3 Antibody



PACO40554

---

## Product Information

**Size:**

50ug

**Reactivity:**

Human

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB, IHC

**Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:1000-1:5000,  
IHC:1:20-1:200

**Protein Background:**

PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. May be involved in pre-mRNA splicing.

**Gene ID:**

PPIL3

**Uniprot**

Q9H2H8

**Synonyms:**

Peptidyl-prolyl cis-trans isomerase-like 3 (PPIase) (EC 5.2.1.8) (Cyclophilin J) (CyPJ)  
(Cyclophilin-like protein PPIL3) (Rotamase PPIL3), PPIL3

**Immunogen:**


Recombinant Human Peptidyl-prolyl cis-trans isomerase-like 3 protein (2-161AA).

**Storage:**


Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

## Product Images

---



Western blot. All lanes: PPIL3 antibody at 10 $\mu$ g/ml. Lane 1: U87 whole cell lysate. Lane 2: HL60 whole cell lysate. Lane 3: K562 whole cell lysate. Lane 4: THP-1 whole cell lysate. Lane 5: A549 whole cell lysate. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 19 kDa. Observed band size: 19 kDa.



Immunohistochemistry of paraffin-embedded human small intestine tissue using PACO40554 at dilution of 1:100.