FANCD2 Antibody, FITC conjugated

PACO40744



Product Information	
Size:	Protein Background:
50ug	Required for maintenance of chromosomal stability. Promotes accurate and efficient
Reactivity:	pairing of homologs during meiosis. Involved in the repair of DNA double-strand breaks, both by homologous recombination and single-strand annealing. May
Human	participate in S phase and G2 phase checkpoint activation upon DNA damage. Plays a role in preventing breakage and loss of missegregating chromatin at the end of cell
Source:	division, particularly after replication stress. Required for the targeting, or stabilization,
Rabbit	of BLM to non-centromeric abnormal structures induced by replicative stress. Promotes BRCA2/FANCD1 loading onto damaged chromatin. May also be involved in B-cell
lsotype:	immunoglobulin isotype switching.
lgG	Gene ID:
Applications:	FANCD2
ELISA	Uniprot
Recommended dilutions:	Q9BXW9
	Synonyms:
	Fanconi anemia group D2 protein (Protein FACD2), FANCD2, FACD
	Immunogen:
	Recombinant Human Fanconi anemia group D2 protein (1-291AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A