## POLR3C Antibody, FITC conjugated

## PACO40972



Product Information	
Size:	Protein Background:
50ug	DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using
Reactivity:	the four ribonucleoside triphosphates as substrates. Specific core component of RNA polymerase III which synthesizes small RNAs, such as 5S rRNA and tRNAs. May direct
Human	with other members of the subcomplex RNA Pol III binding to the TFIIIB-DNA complex via the interactions between TFIIIB and POLR3F. May be involved either in the
Source:	recruitment and stabilization of the subcomplex within RNA polymerase III, or in stimulating catalytic functions of other subunits during initiation. Plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts, such as Epstein-Barr virus-encoded RNAs (EBERs) induce type
Rabbit	
lsotype:	
lgG	
Applications:	I interferon and NF- Kappa-B through the RIG-I pathway.
ELISA	Gene ID:
Recommended dilutions:	POLR3C
	Uniprot
	Q9BUI4
	Synonyms:
	DNA-directed RNA polymerase III subunit RPC3 (RNA polymerase III subunit C3) (DNA- directed RNA polymerase III subunit C) (RNA polymerase III 62 kDa subunit) (RPC62), POLR3C
	Immunogen:
	Recombinant Human DNA-directed RNA polymerase III subunit RPC3 protein (1- 534AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A