NANOG Antibody, Biotin conjugated

PACO42197



Product Information	
Size:	Protein Background:
50ug	Transcription regulator involved in inner cell mass and embryonic stem (ES) cells
Reactivity:	proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks
Human	bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators
Source:	to the active SMAD transcriptional complexes. Acts as a transcriptional activator or
Rabbit	repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'- [CG][GA][CG]C[GC]ATTAN[GC]-3'. Able to autorepress its expression in differentiating
lsotype:	(ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When
lgG	overexpressed, promotes cells to enter into S phase and proliferation.
Applications:	Gene ID:
ELISA	NANOG
Recommended dilutions:	Uniprot
	Q9H9S0
	Synonyms:
	Homeobox protein NANOG (Homeobox transcription factor Nanog) (hNanog), NANOG
	Immunogen:
	Recombinant Human Homeobox protein NANOG protein (1-305AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A