

PACO42262

Product Information

Size:

50ug

Reactivity:

Human, Rat

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC, IF

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:500-1:5000,
IHC:1:20-1:200, IF:1:50-1:200

Protein Background:

Plays a role as an inducible effector molecule that mediates Fas resistance produced by surface Ig engagement in B cells.

Gene ID:

FAIM

Uniprot

Q9NVQ4

Synonyms:

Fas apoptotic inhibitory molecule 1, FAIM, FAIM1


Immunogen:

Recombinant Human Fas apoptotic inhibitory molecule 1 protein (1-179AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4


Product Images



Western blot. All lanes: FAIM antibody at 2 μ g/ml. Lane 1: 293T whole cell lysate. Lane 2: Rat gonad tissue. Lane 3: Jurkat whole cell lysate. Secondary: Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 21, 25, 23 kDa. Observed band size: 21 kDa.



Immunofluorescence staining of HepG2 cells with PACO42262 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemistry of paraffin-embedded human breast cancer using PACO42262 at dilution of 1:100.