

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200

Protein Background:

Oxidoreductase required for tumor suppression. NADPH-bound form inhibits nuclear import by competing with nuclear import substrates for binding to a subset of nuclear transport receptors. May act as a redox sensor linked to transcription through regulation of nuclear import. Isoform 1 is a metastasis suppressor with proapoptotic as well as antiangiogenic properties. Isoform 2 has an antiapoptotic effect.

Gene ID:

HTATIP2

Uniprot

Q9BUP3

Synonyms:

Oxidoreductase HTATIP2 (EC 1.1.1. -) (30 kDa HIV-1 TAT-interacting protein) (HIV-1 TAT-interactive protein 2), HTATIP2, CC3 TIP30


Immunogen:

Recombinant Human Oxidoreductase HTATIP2 protein (1-230AA).


Storage:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Product Images



Immunohistochemistry of paraffin-embedded human small intestine tissue using PACO43198 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human breast cancer using PACO43198 at dilution of 1:100.