

TNFSF13B Antibody



PACO43559

Product Information

Size:

50ul

Reactivity:

Human, Mouse

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IHC

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:500-1:2000,
IHC:1:20-1:200

Protein Background:

Cytokine that binds to TNFRSF13B/TACI and TNFRSF17/BCMA. TNFSF13/APRIL binds to the same 2 receptors. Together, they form a 2 ligands -2 receptors pathway involved in the stimulation of B- and T-cell function and the regulation of humoral immunity. A third B-cell specific BAFF-receptor (BAFFR/BR3) promotes the survival of mature B-cells and the B-cell response.

Gene ID:

TNFSF13B

Uniprot

Q9Y275

Synonyms:

Tumor necrosis factor ligand superfamily member 13B (B lymphocyte stimulator) (BLyS) (B-cell-activating factor) (BAFF) (Dendritic cell-derived TNF-like molecule) (TNF- and APOL-related leukocyte expressed ligand 1) (TALL-1) (CD antigen CD257) [Cleaved into: Tumor necrosis factor ligand superfamily member 13b, membrane form; Tumor necrosis factor ligand superfamily member 13b, soluble form], TNFSF13B, BAFF BLYS TALL1 TNFSF20 ZTNF4


Immunogen:

Recombinant Human Tumor necrosis factor ligand superfamily member 13B protein (76-285AA).


Storage:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Product Images



Western blot. All lanes: Tumor necrosis factor ligand superfamily member 13B antibody at 2 μ g/ml + Mouse liver tissue. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 32, 30, 18 kDa. Observed band size: 32 kDa.



Immunohistochemistry of paraffin-embedded human placenta tissue using PACO43559 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human tonsil tissue using PACO43559 at dilution of 1:100.