

### Product Information

**Size:**

50ul

**Reactivity:**

Human

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, IHC

**Recommended dilutions:**

ELISA:1:2000-1:10000, IHC:1:20-1:200

**Protein Background:**

Dual specificity kinase. Is activated by cytokines and environmental stress in vivo. Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in the MAP kinase p38.

**Gene ID:**

MAP2K3

**Uniprot**

P46734

**Synonyms:**

Dual specificity mitogen-activated protein kinase kinase 3 (MAP kinase kinase 3) (MAPKK 3) (EC 2.7.12.2) (MAPK/ERK kinase 3) (MEK 3) (Stress-activated protein kinase kinase 2) (SAPK kinase 2) (SAPKK-2) (SAPKK2), MAP2K3, MEK3 MKK3 PRKMK3 SKK2

**Immunogen:**


Recombinant Human Dual specificity mitogen-activated protein kinase kinase 3 protein (1-170AA).

**Storage:**


PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Product Images


---



Immunohistochemistry of paraffin-embedded human endometrial cancer using PACO43815 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human placenta tissue using PACO43815 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human tonsil tissue using PACO43815 at dilution of 1:100.