

NMT2 Antibody



PACO44511

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC, IP

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200,
IP:1:200-1:2000

Protein Background:

Adds a myristoyl group to the N-terminal glycine residue of certain cellular and viral proteins.

Gene ID:

NMT2

Uniprot

O60551

Synonyms:

Glycylpeptide N-tetradecanoyltransferase 2 (EC 2.3.1.97) (Myristoyl-CoA: protein N-myristoyltransferase 2) (NMT 2) (Peptide N-myristoyltransferase 2) (Type II N-myristoyltransferase), NMT2


Immunogen:

Recombinant Human Glycylpeptide N-tetradecanoyltransferase 2 protein (1-120AA).


Storage:

PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Product Images



Immunohistochemistry of paraffin-embedded human testis tissue using PACO44511 at dilution of 1:100.



Immunoprecipitating NMT2 in HepG2 whole cell lysate. Lane 1: Rabbit control IgG instead of (1 μ g) instead of PACO44511 in HepG2 whole cell lysate. For western blotting, a HRP-conjugated anti-rabbit IgG, specific to the non-reduced form of IgG was used as the Secondary antibody (1/50000). Lane 2: PACO44511 (4 μ g) + HepG2 whole cell lysate (500 μ g). Lane 3: HepG2 whole cell lysate (20 μ g).



Immunohistochemistry of paraffin-embedded human prostate cancer using PACO44511 at dilution of 1:100.