

PACO44553

---

## Product Information

**Size:**

50ul

**Reactivity:**

Human, Mouse

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB, IHC

**Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:1000-1:5000,  
IHC:1:20-1:200

**Protein Background:**

Acts as a negative mediator of apoptosis and may play a role in tumor progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.

**Gene ID:**

TNFAIP8

**Uniprot**

O95379

**Synonyms:**

Tumor necrosis factor alpha-induced protein 8 (TNF alpha-induced protein 8) (Head and neck tumor and metastasis-related protein) (MDC-3.13) (NF-kappa-B-inducible DED-containing protein) (NDED) (SCC-S2) (TNF-induced protein GG2-1), TNFAIP8

**Immunogen:**


Recombinant Human Tumor necrosis factor alpha-induced protein 8 protein (1-198AA).

**Storage:**


PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Product Images

---



Western blot. All lanes: TNFAIP8 antibody at 2.79 $\mu$ g/ml. Lane 1: A431 whole cell lysate. Lane 2: PC-3 whole cell lysate. Lane 3: Human placenta tissue. Lane 4: Mouse thymus tissue. Secondary: Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 24, 23, 22, 25 kDa. Observed band size: 22 kDa.



Immunohistochemistry of paraffin-embedded human colon cancer using PACO44553 at dilution of 1:100.