

# USP26 Antibody



PACO44809

---

## Product Information

**Size:**

50ul

**Reactivity:**

Human, Mouse

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB, IHC

**Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:1000-1:5000,  
IHC:1:20-1:200

**Protein Background:**

Involved in the ubiquitin-dependent proteolytic pathway in conjunction with the 26S proteasome. Deubiquitinates the androgen receptor and regulates the androgen receptor signaling pathway.

**Gene ID:**

USP26

**Uniprot**

Q9BXU7

**Synonyms:**

Ubiquitin carboxyl-terminal hydrolase 26 (EC 3.4.19.12) (Deubiquitinating enzyme 26) (Ubiquitin thioesterase 26) (Ubiquitin-specific-processing protease 26), USP26

**Immunogen:**


Recombinant Human Ubiquitin carboxyl-terminal hydrolase 26 protein (664-913AA).

**Storage:**


PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Product Images


---



Western blot. All lanes: USP26 antibody at 5.22 $\mu$ g/ml. Lane 1: HeLa whole cell lysate. Lane 2: U937 whole cell lysate. Lane 3: Mouse gonadal tissue. Lane 4: Mouse liver tissue. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 105 kDa. Observed band size: 105 kDa.



Immunohistochemistry of paraffin-embedded human small intestine tissue using PACO44809 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human liver tissue using PACO44809 at dilution of 1:100.