## **AHR Antibody, Biotin conjugated**



## PACO45657

Rabbit

## **Product Information**

Size: Protein Background:

50ug Ligand-activated transcriptional activator. Binds to the XRE promoter region of genes it

**Reactivity:** activates. Activates the expression of multiple phase I and II xenobiotic chemical metabolizing enzyme genes (such as the CYP1A1 gene). Mediates biochemical and

Human toxic effects of halogenated aromatic hydrocarbons. Involved in cell-cycle regulation.

Likely to play an important role in the development and maturation of many tissues.

Source: Regulates the circadian clock by inhibiting the basal and circadian expression of the

core circadian component PER1. Inhibits PER1 by repressing the CLOCK-ARNTL/BMAL1

Core circadian component reixt. Inhibits reixt by repressing the CLOCK-AKIVILY BIVIA

heterodimer mediated transcriptional activation of PER1.

Isotype: Gene ID:

IgG AHR

Applications: Uniprot

ELISA P35869

Recommended dilutions: Synonyms:

Aryl hydrocarbon receptor (Ah receptor) (AhR) (Class E basic helix-loop-helix protein 76)

(bHLHe76), AHR, BHLHE76

Immunogen:

Recombinant Human Aryl hydrocarbon receptor protein (126-353AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product	<b>Images</b>
---------	---------------

N/A N/A