
## HIST1H4A Antibody

## PACO45902




Product Information	
Size:	Protein Background:
50ug	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Reactivity:	
Human, Mouse	
Source:	
Rabbit	Gene ID:
lsotype:	HIST1H4A
lgG	Uniprot
Applications:	P62805
ELISA, WB, IHC	Synonyms:
Recommended dilutions:	Histone H4, HIST1H4A; HIST1H4B; HIST1H4C; HIST1H4D; HIST1H4E; HIST1H4F; HIST1H4H; HIST1H4I; HIST1H4J; HIST1H4K; HIST1H4L; HIST2H4A; HIST2H4B; HIST4H4, H4/A H4FA; H4/I H4FI; H4/G H4FG; H4/B H4FB; H4/J H4FJ; H4/C H4FC; H4/H H4FH; H4/M H4FM; H4/E H4FE; H4/D H4FD; H4/K H4FK; H4/N H4F2 H4FN HIST2H4; H4/O H4FO;
ELISA:1:2000-1:10000, WB:1:1000-1:5000, IHC:1:20-1:200	
	Immunogen:
	Recombinant Human Histone H4 protein (2-50AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4



Western blot. All lanes: HIST1H4A antibody at 14  $\mu$ g/ml + Mouse thymus tissue. Secondary. Goat polyclonal to rabbit IgG at 1/10000 dilution. Predicted band size: 11 kDa. Observed band size: 11 kDa.



Immunohistochemistry of paraffin-embedded human liver cancer using PACO45902 at dilution of 1:100.