ITGA4 Antibody, HRP conjugated



PACO45955

lgG

Product Information

Size: Protein Background:

50ug Integrins alpha-4/beta-1 (VLA-4) and alpha-4/beta-7 are receptors for fibronectin. They recognize one or more domains within the alternatively spliced CS-1 and CS-5 regions

Reactivity:of fibronectin. They are also receptors for VCAM1. Integrin alpha-4/beta-1 recognizes

Human the sequence Q-I-D-S in VCAM1. Integrin alpha-4/beta-7 is also a receptor for

MADCAM1. It recognizes the sequence L-D-T in MADCAM1. On activated endothelial **Source:**

cells integrin VLA-4 triggers homotypic aggregation for most VLA-4-positive leukocyte

Rabbit cell lines. It may also participate in cytolytic T-cell interactions with target cells. ITGA4:

ITGB1 binds to fractalkine (CX3CL1) and may act as its coreceptor in CX3CR1-

Isotype: dependent fractalkine signaling. ITGA4: ITGB1 binds to PLA2G2A via a site (site 2) which

is distinct from the classical ligand-binding site (site 1) and this induces integrin

conformational changes and enhanced ligand binding to site 1.

Applications: Gene ID:

ELISA ITGA4

Recommended dilutions: Uniprot

P13612

Synonyms:

Integrin alpha-4 (CD49 antigen-like family member D) (Integrin alpha-IV) (VLA-4 $\,$

subunit alpha) (CD antigen CD49d), ITGA4, CD49D

Immunogen:

Recombinant Human Integrin alpha-4 protein (376-558AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product	Images
---------	---------------

N/A N/A