

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200

Protein Background:

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Gene ID:

MED31

Uniprot

Q9Y3C7

Synonyms:

Mediator of RNA polymerase II transcription subunit 31 (Mediator complex subunit 31) (Mediator complex subunit SOH1) (hSOH1), MED31, SOH1


Immunogen:

Recombinant Human Mediator of RNA polymerase II transcription subunit 31 protein (2-72AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product Images



Immunohistochemistry of paraffin-embedded human pancreatic tissue using PACO46022 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human small intestine tissue using PACO46022 at dilution of 1:100.