PAN2 Antibody, HRP conjugated

PACO46583



Product Information	
Size:	Protein Background:
50ug	Catalytic subunit of the poly(A)-nuclease (PAN) deadenylation complex, one of two
Reactivity:	cytoplasmic mRNA deadenylases involved in general and miRNA-mediated mRNA turnover. PAN specifically shortens poly(A) tails of RNA when the poly(A) stretch is
Human	bound by poly(A)-binding protein (PABP), which is followed by rapid degradation of the shortened mRNA tails by the CCR4-NOT complex. Deadenylated mRNAs are then
Source:	degraded by two alternative mechanisms, namely exosome-mediated 3'-5'
Rabbit	exonucleolytic degradation, or deadenlyation-dependent mRNA decaping and subsequent 5'-3' exonucleolytic degradation by XRN1.
lsotype:	Gene ID:
lgG	PAN2
Applications:	Uniprot
ELISA	Q504Q3
Recommended dilutions:	Synonyms:
	PAN2-PAN3 deadenylation complex catalytic subunit PAN2 (EC 3.1.13.4) (Inactive ubiquitin carboxyl-terminal hydrolase 52) (PAB1P-dependent poly(A)-specific ribonuclease) (Poly(A)-nuclease deadenylation complex subunit 2) (PAN deadenylation complex subunit 2), PAN2, KIAA0710 USP52
	Immunogen:
	Recombinant Human PAB-dependent poly(A)-specific ribonuclease subunit PAN2 protein (628-908AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A