

# MED25 Antibody, FITC conjugated



PACO46652

---

## Product Information

**Size:**

50ug

**Protein Background:**

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. Required for RARA/RXRA-mediated transcription.

**Reactivity:**

Human

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA

**Recommended dilutions:****Gene ID:**

MED25

**Uniprot**

Q71SY5

**Synonyms:**

Mediator of RNA polymerase II transcription subunit 25 (Activator interaction domain-containing protein 1) (Activator-recruited cofactor 92 kDa component) (ARC92) (Mediator complex subunit 25) (p78), MED25, acid, ARC92 PTOV2

**Immunogen:**

Recombinant Human Mediator of RNA polymerase II transcription subunit 25 protein (112-140AA).

**Storage:**

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

## Product Images

---

N/A

N/A