

---

## Product Information

**Size:**

50ug

**Protein Background:**

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA

**Reactivity:**

Human

polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

**Source:**

Rabbit

**Gene ID:**

MED17

**Isotype:**

IgG

**Uniprot****Applications:**

ELISA

**Q9NVC6****Recommended dilutions:**

Mediator of RNA polymerase II transcription subunit 17 (Activator-recruited cofactor 77 kDa component) (ARC77) (Cofactor required for Sp1 transcriptional activation subunit 6) (CRSP complex subunit 6) (Mediator complex subunit 17) (Thyroid hormone receptor-associated protein complex 80 kDa component) (Trap80) (Transcriptional coactivator CRSP77) (Vitamin D3 receptor-interacting protein complex 80 kDa component) (DRIP80), MED17, ARC77 CRSP6 DRIP77 DRIP80 TRAP80

**Immunogen:**

Recombinant Human Mediator of RNA polymerase II transcription subunit 17 protein (411-651AA).

**Storage:**

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

## Product Images

---

N/A

N/A