

MED17 Antibody, FITC conjugated



PACO48104

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA

Recommended dilutions:**Protein Background:**

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Gene ID:

MED17

Uniprot

Q9NVC6

Synonyms:

Mediator of RNA polymerase II transcription subunit 17 (Activator-recruited cofactor 77 kDa component) (ARC77) (Cofactor required for Sp1 transcriptional activation subunit 6) (CRSP complex subunit 6) (Mediator complex subunit 17) (Thyroid hormone receptor-associated protein complex 80 kDa component) (Trap80) (Transcriptional coactivator CRSP77) (Vitamin D3 receptor-interacting protein complex 80 kDa component) (DRIP80), MED17, ARC77 CRSP6 DRIP77 DRIP80 TRAP80

Immunogen:

Recombinant Human Mediator of RNA polymerase II transcription subunit 17 protein (411-651AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Product Images

N/A

N/A