## RACGAP1 Antibody, Biotin conjugated

## PACO48149



| Product Information    |   |
|------------------------|---|
| Size:                  | Protein Background:   |
| 50ug                   | Component of the centralspindlin complex that serves as a microtubule-dependent and   |
| Reactivity:            | Rho-mediated signaling required for the myosin contractile ring formation during the<br>cell cycle cytokinesis. Required for proper attachment of the midbody to the cell     |
| Human                  | membrane during cytokinesis. Plays key roles in controlling cell growth and differentiation of hematopoietic cells through mechanisms other than regulating Rac               |
| Source:                | GTPase activity. Also involved in the regulation of growth-related processes in   |
| Rabbit                 | adipocytes and myoblasts. May be involved in regulating spermatogenesis and in the RACGAP1 pathway in neuronal proliferation. Shows strong GAP (GTPase activation)            |
| lsotype:               | activity towards CDC42 and RAC1 and less towards RHOA. Essential for the early stages   |
| lgG                    | of embryogenesis. May play a role in regulating cortical activity through RHOA during cytokinesis. May participate in the regulation of sulfate transport in male germ cells. |
| Applications:          | Gene ID:  |
| ELISA                  | RACGAP1   |
| Recommended dilutions: | Uniprot   |
|                        | Q9H0H5  |
|                        | Synonyms:   |
|                        | Rac GTPase-activating protein 1 (Male germ cell RacGap) (MgcRacGAP) (Protein CYK4<br>homolog) (CYK4) (HsCYK-4), RACGAP1, KIAA1478 MGCRACGAP                                   |
|                        | Immunogen:  |
|                        | Recombinant Human Rac GTPase-activating protein 1 protein (86-331AA).   |
|                        | Storage:  |
|                        | Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  |

N/A N/A