CCND2 Antibody, Biotin conjugated

PACO48861



Product Information	
Size:	Protein Background:
50ug	Regulatory component of the cyclin D2-CDK4 (DC) complex that phosphorylates and
Reactivity:	inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the
Human	transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase.
Source:	Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major
Rabbit	integrators of various mitogenenic and antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its
lsotype:	transcriptional activity. Component of the ternary complex, cyclin D2/CDK4/CDKN1B,
lgG	required for nuclear translocation and activity of the cyclin D-CDK4 complex.
Applications:	CCND2
ELISA	Uniprot
Recommended dilutions:	P30279
	Synonyms:
	G1/S-specific cyclin-D2, CCND2
	Immunogen:
	Recombinant Human G1/S-specific cyclin-D2 protein (215-285AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

N/A N/A