KLRK1 Antibody, FITC conjugated

PACO48992



| Product Information | |
|------------------------|--|
| Size: | Protein Background: |
| 50ug | Function as an activating and costimulatory receptor involved in immunosurveillance |
| Reactivity: | upon binding to various cellular stress-inducible ligands displayed at the surface of autologous tumor cells and virus-infected cells. Provides both stimulatory and |
| Human | costimulatory innate immune responses on activated killer (NK) cells, leading to cytotoxic activity. Acts as a costimulatory receptor for T-cell receptor (TCR) in CD8(+) T- |
| Source: | cell-mediated adaptive immune responses by amplifying T-cell activation. Stimulates perforin-mediated elimination of ligand-expressing tumor cells. Signaling involves calcium influx, culminating in the expression of TNF-alpha. Participates in NK cell- mediated bone marrow graft rejection. May play a regulatory role in differentiation and survival of NK cells. Binds to ligands belonging to various subfamilies of MHC class I- related glycoproteins including MICA, MICB, RAET1E, RAET1G, ULBP1, ULBP2, ULBP3 (ULBP2>ULBP1>ULBP3) and ULBP4. Gene ID: |
| Rabbit | |
| lsotype: | |
| lgG | |
| Applications: | |
| ELISA | |
| Recommended dilutions: | KLRK1 |
| | Uniprot |
| | P26718 |
| | Synonyms: |
| | NKG2-D type II integral membrane protein (Killer cell lectin-like receptor subfamily K member 1) (NK cell receptor D) (NKG2-D-activating NK receptor) (CD antigen CD314), KLRK1, D12S2489E NKG2D |
| | Immunogen: |
| | Recombinant Human NKG2-D type II integral membrane protein (73-216AA). |
| | Storage: |
| | Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |

N/A N/A