RNF40 Antibody, FITC conjugated



PACO49960

Source:

Product Information

Size: Protein Background:

50ug Component of the RNF20/40 E3 ubiquitin-protein ligase complex that mediates monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1). H2BK120ub1 gives a

Reactivity: specific tag for epigenetic transcriptional activation and is also prerequisite for histone

Human H3 'Lys-4' and 'Lys-79' methylation (H3K4me and H3K79me, respectively). It thereby

plays a central role in histone code and gene regulation. The RNF20/40 complex forms a H2B ubiquitin ligase complex in cooperation with the E2 enzyme UBE2A or UBE2B;

reports about the cooperation with UBE2E1/UBCH are contradictory. Required for

Rabbit transcriptional activation of Hox genes.

Isotype: Gene ID:

IgG RNF40

Applications: Uniprot

ELISA 075150

Recommended dilutions: Synonyms:

E3 ubiquitin-protein ligase BRE1B (BRE1-B) (EC 2.3.2.27) (95 kDa retinoblastoma-associated protein) (RBP95) (RING finger protein 40) (RING-type E3 ubiquitin

transferase BRE1B), RNF40, BRE1B KIAA0661

Immunogen:

Recombinant Human E3 ubiquitin-protein ligase BRE1B protein (471-724AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product	Images
---------	---------------

N/A N/A