## NOD2 Antibody, HRP conjugated

## PACO51095



Product Information	
Size:	Protein Background:
50ug	Involved in gastrointestinal immunity. Upon stimulation by muramyl dipeptide (MDP), a
Reactivity:	fragment of bacterial peptidoglycan, binds the proximal adapter receptor-interacting RIPK2, which recruits ubiquitin ligases as XIAP, BIRC2, BIRC3 and the LUBAC complex,
Human	triggering activation of MAP kinases and activation of NF-kappa-B signaling. This in turn leads to the transcriptional activation of hundreds of genes involved in immune
Source:	response. Required for MDP-induced NLRP1-dependent CASP1 activation and IL1B
Rabbit	release in macrophages.
lsotype:	Gene ID:
lgG	NOD2
Applications:	Uniprot
ELISA	Q9HC29
Recommended dilutions:	Synonyms:
	Nucleotide-binding oligomerization domain-containing protein 2 (Caspase recruitment domain-containing protein 15) (Inflammatory bowel disease protein 1), NOD2, CARD15 IBD1
	Immunogen:
	Recombinant Human Nucleotide-binding oligomerization domain-containing protein 2 protein (460-733AA).
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

N/A N/A