EIF3A Antibody, HRP conjugated



PACO52399

Reactivity:

Source:

Rabbit

Isotype:

lgG

Product Information

Size: Protein Background:

50ug Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex

associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2: GTP: methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The

Human

GTP: methionyi-tkiNAI and eIF-5 to form the 435 pre-initiation complex (435 PIC). The

eIF-3 complex stimulates mRNA recruitment to the 435 PIC and scanning of the mRNA

for AUG recognition. The eIF-3 complex is also required for disassembly and recycling

of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Essential for the initiation of

translation on type-1 viral ribosomal entry sites (IRESs), like for HCV, PV, EV71 or BEV

translation. In case of FCV infection, plays a role in the ribosomal termination-

reinitiation event leading to the translation of VP2.

Applications: Gene ID:

ELISA EIF3A

Recommended dilutions: Uniprot

Q14152

Synonyms:

Eukaryotic translation initiation factor 3 subunit A (eIF3a) (Eukaryotic translation initiation factor 3 subunit 10) (eIF-3-theta) (eIF3 p167) (eIF3 p180) (eIF3 p185), EIF3A,

EIF3S10 KIAA0139

Immunogen:

Recombinant Human Eukaryotic translation initiation factor 3 subunit A protein (1288-

1379AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product	Images
---------	---------------

N/A N/A