CD207 Antibody, Biotin conjugated

PACO55365



Product Information	
Size:	Protein Background:
50ug	Calcium-dependent lectin displaying mannose-binding specificity. Induces the
Reactivity:	formation of Birbeck granules (BGs); is a potent regulator of membrane superimposition and zippering. Binds to sulfated as well as mannosylated glycans,
Human	keratan sulfate (KS) and beta-glucans. Facilitates uptake of antigens and is involved in the routing and/or processing of antigen for presentation to T cells. Major receptor on
Source:	primary Langerhans cells for Candida species, Saccharomyces species, and Malassezia
Rabbit	furfur. Protects against human immunodeficiency virus-1 (HIV-1) infection. Binds to high-mannose structures present on the envelope glycoprotein which is followed by
lsotype:	subsequent targeting of the virus to the Birbeck granules leading to its rapid degradation.
lgG	Gene ID:
Applications:	CD207
ELISA	Uniprot
Recommended dilutions:	Q9UJ71
	Synonyms:
	C-type lectin domain family 4 member K (Langerin) (CD_antigen: CD207), CD207, CLEC4K
	Immunogen:
	Recombinant Human C-type lectin domain family 4 member K protein (229-312AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

N/A N/A