MASTL Antibody, FITC conjugated

PACO56068



Product Information	
Size:	Protein Background:
50ug	Serine/threonine kinase that plays a key role in M phase by acting as a regulator of
Reactivity:	mitosis entry and maintenance. Acts by promoting the inactivation of protein phosphatase 2A (PP2A) during M phase: does not directly inhibit PP2A but acts by
Human	mediating phosphorylation and subsequent activation of ARPP19 and ENSA at 'Ser-62' and 'Ser-67', respectively. ARPP19 and ENSA are phosphatase inhibitors that specifically
Source:	inhibit the PPP2R2D (PR55-delta) subunit of PP2A. Inactivation of PP2A during M phase
Rabbit	is essential to keep cyclin-B1-CDK1 activity high. Following DNA damage, it is also involved in checkpoint recovery by being inhibited. Phosphorylates histone protein in
lsotype:	vitro; however such activity is unsure in vivo. May be involved in megakaryocyte differentiation.
lgG	Gene ID:
Applications:	MASTL
ELISA	Uniprot
Recommended dilutions:	Q96GX5
	Synonyms:
	Serine/threonine-protein kinase greatwall (GW) (GWL) (hGWL) (EC 2.7.11.1) (Microtubule-associated serine/threonine-protein kinase-like) (MAST-L), MASTL, GW GWL THC2
	Immunogen:
	Recombinant Human Serine/threonine-protein kinase greatwall protein (612-740AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

N/A N/A