
Di-methyl-HIST1H3A (K36) Antibody

PACO56505



Product Information	
Size:	Protein Background:
50ul	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Reactivity:	
Human	
Source:	
Rabbit	Gene ID:
lsotype:	HIST1H3A
lgG	Uniprot
Applications:	P68431
ELISA, IF	Synonyms:
Recommended dilutions:	Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ
ELISA:1:2000-1:10000, IF:1:50-1:200	
	Immunogen:
	Peptide sequence around site of Di-methyl-Lys (36) derived from Human Histone H3.1.
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



Immunofluorescent analysis of Hela cells using PACO56505 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).