

# HIST1H3A (Ab-4) Antibody



PACO56531

---

## Product Information

**Size:**

50ul

**Reactivity:**

Human, Mouse

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB, IHC, IF, CHIP

**Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:100-1:1000,  
IHC:1:20-1:200, IF:1:50-1:200

**Protein Background:**

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

**Gene ID:**

HIST1H3A

**Uniprot**

P68431

**Synonyms:**

Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ

**Immunogen:**


Peptide sequence around site of Lys (4) derived from Human Histone H3.1.

**Storage:**


Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## Product Images


---



Western Blot. Positive WB detected in: NIH/3T3 cell acid extracts, HeLa cell acid extracts, A549 cell acid extracts, PC-3 cell acid extracts, HEK293 cell acid extracts, Mouse kidney tissue, Mouse brain tissue. All lanes: HIST1H3A antibody at 2.5µg/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 16 kDa. Observed band size: 16 kDa.



Immunohistochemistry of paraffin-embedded human lung cancer using PACO56531 at dilution of 1:100.



Immunohistochemistry of paraffin-embedded human colon cancer using PACO56531 at dilution of 1:100.