## Acetyl-HIST1H3A (K14) Antibody



## PACO56564

Reactivity:

Human

## **Product Information**

Size: Protein Background:

50ul Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template.

Histones thereby play a central role in transcription regulation, DNA repair, DNA

replication and chromosomal stability. DNA accessibility is regulated via a complex set

of post-translational modifications of histones, also called histone code, and

**Source:** nucleosome remodeling.

Rabbit Gene ID:

Isotype: HIST1H3A

lgG **Uniprot** 

**Applications:** P68431

ELISA:1:2000-1:10000, IF:1:50-1:200

ELISA, IF Synonyms:

Recommended dilutions: Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/d)

H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK;

H3FF; H3FJ


Immunogen:

Peptide sequence around site of Acetyl-Lys (14) derived from Human Histone H3.1.

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## **Product Images**



Immunofluorescent analysis of Hela cells treated by NaB using PACO56564 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).