HIST1H3A (Ab-9) Antibody



PACO56574

Reactivity:

Source:

Product Information

Size: Protein Background:

50ul Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template.

Histones thereby play a central role in transcription regulation, DNA repair, DNA

Human, Mouse replication and chromosomal stability. DNA accessibility is regulated via a complex set

of post-translational modifications of histones, also called histone code, and

nucleosome remodeling.

Rabbit Gene ID:

Isotype: HIST1H3A

lgG **Uniprot**

Applications: P68431

ELISA, WB, IF, ChIP Synonyms:

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:100-1:1000, IF:1:50-1:200

Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ


Immunogen:


Peptide sequence around site of Lys (9) derived from Human Histone H3.1.


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images







Chromatin Immunoprecipitation Hela (4*10^6

) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with $8\mu g$ anti-HIST1H3A (PACO56574) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the beta -Globin promoter.

Immunofluorescent analysis of Hela cells using PACO56574 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IqG(H+L).