

Phospho-HIST1H1B (S17) Antibody



PACO56605

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, ICC, IF

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:100-1:1000,
ICC:1:20-1:200, IF:1:50-1:200

Protein Background:

Histone H1 protein binds to linker DNA between nucleosomes forming the macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation.

Gene ID:

HIST1H1B

Uniprot

P16401

Synonyms:

Histone H1.5 (Histone H1a) (Histone H1b) (Histone H1s-3), HIST1H1B, H1F5


Immunogen:

Peptide sequence around site of Phospho-Ser (17) derived from Human Histone H1.5.


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4


Product Images



Western Blot. Positive WB detected in: MCF-7 whole cell lysate. All lanes: HIST1H1B antibody at 1 μ g/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 23 kDa. Observed band size: 32 kDa.



Immunofluorescence staining of HepG2 cells with PACO56605 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunocytochemistry analysis of HepG2 cells using PACO56605 at dilution of 1:100.