
HIST1H1E (Ab-17) Antibody

PACO56694



Product Information	
Size:	Protein Background:
50ul	Histone H1 protein binds to linker DNA between nucleosomes forming the
Reactivity:	macromolecular structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation. Gene ID:
Human, Rat	
Source:	
Rabbit	HIST1H1E
lsotype:	Uniprot
lgG	P10412
Applications:	Synonyms:
ELISA, WB, IHC, IF	Histone H1.4 (Histone H1b) (Histone H1s-4), HIST1H1E, H1F4
Recommended dilutions:	Immunogen:
ELISA:1:2000-1:10000, WB:1:50-1:500, IHC:1:20-1:200, IF:1:1-1:10	Peptide sequence around site of Thr (17) derived from Human Histone H1.4.
	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



Western Blot. Positive WB detected in: Hela whole cell lysate, K562 whole cell lysate, Rat brain tissue. All lanes: HIST1H1E antibody at 1.48μ g/ml. Secondary. Goat polyclonal to rabbit lgG at 1/50000 dilution. Predicted band size: 22 kDa. Observed band size: 28 kDa.

Immunofluorescence staining of Hela cells with PACO56694 at 1:2.5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

IHC image of PACO56694 diluted at 1:50 and staining in paraffinembedded human glioma performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.