GATAD2A Antibody



PACO57284

Product Information

Size: **Protein Background:**

50ug Transcriptional repressor. Enhances MBD2-mediated repression. Efficient repression

requires the presence of GATAD2B.

Reactivity: Gene ID:

Human, Rat GATAD2A

Source: Uniprot Rabbit

Q86YP4 Isotype:

Synonyms: lgG

Transcriptional repressor p66-alpha (Hp66alpha) (GATA zinc finger domain-containing **Applications:**

protein 2A), GATAD2A

ELISA, WB, IHC Immunogen:

Recommended dilutions: Recombinant Human Transcriptional repressor p66-α protein (170-241AA).


ELISA:1:2000-1:10000, WB:1:500-1:5000,


IHC:1:500-1:1000


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4


Product Images







IHC image of PACO57284 diluted at 1:800 and staining in paraffinembedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of PACO57284 diluted at 1:800 and staining in paraffinembedded human small intestine tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.