

PACO57596

---

## Product Information

**Size:**

50ug

**Reactivity:**

Human

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, IHC

**Recommended dilutions:**

ELISA:1:2000-1:10000, IHC:1:20-1:200

**Protein Background:**

Involved in RNA polymerase III-mediated transcription. Integral, tightly associated component of the DNA-binding TFIIIC2 subcomplex that directly binds tRNA and virus-associated RNA promoters.

**Gene ID:**

GTF3C5

**Uniprot**

Q9Y5Q8

**Synonyms:**

General transcription factor 3C polypeptide 5 (TF3C-epsilon) (Transcription factor III C 63 kDa subunit) (TFIIIC 63 kDa subunit) (TFIIIC63) (Transcription factor IIC subunit epsilon), GTF3C5

**Immunogen:**


Recombinant Human General transcription factor 3C polypeptide 5 protein (305-448AA).

**Storage:**

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## Product Images

---



IHC image of PACO57596 diluted at 1:100 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.