# HIST1H3A (Ab-18) Antibody



#### PACO57642

Reactivity:

Human

Source:

#### **Product Information**

Size: Protein Background:

50ul Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template.

Histones thereby play a central role in transcription regulation, DNA repair, DNA

replication and chromosomal stability. DNA accessibility is regulated via a complex set

of post-translational modifications of histones, also called histone code, and

nucleosome remodeling.

Rabbit Gene ID:

Isotype: HIST1H3A

lgG Uniprot

**Applications:** P68431

ELISA, IHC, IF Synonyms:

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200, IF:1:10-1:100

Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ


## Immunogen:


Peptide sequence around site of Lys (18) derived from Human Histone H3.1.

### Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4  $\,$ 

## **Product Images**





IHC image of PACO57642 diluted at 1:30 and staining in paraffinembedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of Hela cells with PACO57642 at 1:15, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).