
ADAMTS13 Antibody

PACO57652



| Product Information | |
|---------------------------------------|--|
| Size: | Protein Background: |
| 50ug | Cleaves the vWF multimers in plasma into smaller forms thereby controlling vWF- |
| Reactivity: | mediated platelet thrombus formation. |
| Human | Gene ID: |
| Source: | ADAMTS13 |
| Rabbit | Uniprot |
| lsotype: | Q76LX8 |
| lgG | Synonyms: |
| Applications: | A disintegrin and metalloproteinase with thrombospondin motifs 13 (ADAM-TS 13) (ADAM-TS13) (ADAMTS-13) (EC 3.4.24.87) (von Willebrand factor-cleaving protease) |
| ELISA, IHC | (vWF-CP) (vWF-cleaving protease), ADAMTS13, C9orf8 |
| Recommended dilutions: | Immunogen: |
| ELISA:1:2000-1:10000, IHC:1:200-1:500 | Recombinant Human A disintegrin and metalloproteinase with thrombospondin motifs 13 protein (738-905AA). |
| | Storage: |

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



IHC image of PACO57652 diluted at 1:400 and staining in paraffinembedded human liver tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.