DSCAM Antibody



PACO57736

-				
ш	200		Intori	mation
г	IUU	uct		Hauon

Size: Protein Background:

50ug

Reactivity: Gene ID:

Human DSCAM

Source: Uniprot

Rabbit O60469

Isotype: Synonyms:

IgG Down syndrome cell adhesion molecule (CHD2), DSCAM


Applications: Immunogen:


ELISA, IHC Recombinant Human Down syndrome cell adhesion molecule protein (1711-1867AA).

Recommended dilutions: Storage:

ELISA:1:2000-1:10000, IHC:1:200-1:500 Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images





IHC image of PACO57736 diluted at 1:200 and staining in paraffinembedded human glioma performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

IHC image of PACO57736 diluted at 1:200 and staining in paraffinembedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.