
NKX2-8 Antibody


PACO57892



Product Information	
Size:	Protein Background:
50ug	double-stranded DNA binding, RNA polymerase II core promoter proximal region sequence-specific DNA binding, sequence-specific DNA binding, transcription factor activity, sequence-specific DNA binding, transcriptional activator activity, RNA polymerase II core promoter proximal region sequence-specific binding, liver development, positive regulation of transcription from RNA polymerase II promoter, transcription from RNA polymerase II promoter, transcription, DNA-templated.
Reactivity:	
Human	
Source:	
Rabbit	Gene ID:
lsotype:	NKX2-8
lgG	Uniprot
Applications:	O15522
ELISA, IHC	Synonyms:
Recommended dilutions:	Homeobox protein Nkx-2.8 (Homeobox protein NK-2 homolog H), NKX2-8, NKX-2.8 NKX2G NKX2H
ELISA:1:2000-1:10000, IHC:1:200-1:500	
	Immunogen:
	Recombinant Human Homeobox protein Nkx-2.8 protein (1-160AA).
	Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4





IHC image of PACO57892 diluted at 1:300 and staining in paraffinembedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

IHC image of PACO57892 diluted at 1:300 and staining in paraffinembedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.