
NEB Antibody

PACO58084



Product Information	
Size:	Protein Background:
50ug	This giant muscle protein may be involved in maintaining the structural integrity of sarcomeres and the membrane system associated with the myofibrils. Binds and stabilize F-actin.
Reactivity:	
Human	Gene ID:
Source:	NEB
Rabbit	Uniprot
lsotype:	P20929
lgG	Synonyms:
Applications:	Nebulin, NEB
elisa, IHC, If	Immunogen:
Recommended dilutions:	Recombinant Human Nebulin protein (245-373AA).
ELISA:1:2000-1:10000, IHC:1:500-1:1000, IF:1:200-1:500	Storage:
	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



IHC image of PACO58084 diluted at 1:600 and staining in paraffinembedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of HepG2 cells with PACO58084 at 1:200, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

IHC image of PACO58084 diluted at 1:600 and staining in paraffinembedded human skeletal muscle tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.