LAPTM4B Antibody, Biotin conjugated



PACO58255

Product Information

Size: Protein Background:

50ug Required for optimal lysosomal function. Blocks EGF-stimulated EGFR intraluminal sorting and degradation. Conversely by binding with the phosphatidylinositol 4,5-

Reactivity:bisphosphate, regulates its PIP5K1C interaction, inhibits HGS ubiquitination and relieves

Human LAPTM4B inhibition of EGFR degradation. Recruits SLC3A2 and SLC7A5 (the Leu transporter) to the lysosome, promoting entry of leucine and other essential amino

Source: acid, (EAA) into the lysosome, stimulating activation of proton-transporting vacuolar (V)-ATPase protein pump (V-ATPase) and hence mTORC1 activation. Plays a role as

Rabbit negative regulator of TGFB1 production in regulatory T cells. Binds ceramide and

Isotype: facilitates its exit from late endosome in order to control cell death pathways.

lgG Gene ID:

Applications: LAPTM4B

ELISA Uniprot

Q86VI4 Recommended dilutions:

Synonyms:

Lysosomal-associated transmembrane protein 4B (Lysosome-associated

transmembrane protein 4-beta), LAPTM4B

Immunogen:

Recombinant Human Lysosomal-associated transmembrane protein 4B protein (174-226AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

| Product | Images |
|---------|---------------|
|---------|---------------|

N/A N/A