
## Phospho-HIST1H3A (S28) Antibody

## PACO58653



| Product Information   |   |
|---|---|
| Size:   | Protein Background:   |
| 50ul  | Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin,<br>limiting DNA accessibility to the cellular machineries which require DNA as a template.<br>Histones thereby play a central role in transcription regulation, DNA repair, DNA<br>replication and chromosomal stability. DNA accessibility is regulated via a complex set<br>of post-translational modifications of histones, also called histone code, and<br>nucleosome remodeling. |
| Reactivity:   |   |
| Human   |   |
| Source:   |   |
| Rabbit  | Gene ID:  |
| lsotype:  | HIST1H3A  |
| lgG   | Uniprot   |
| Applications:   | P68431  |
| ELISA, IF, ChIP   | Synonyms:   |
| Recommended dilutions:  | Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone   |
| ELISA:1:2000-1:10000, IF:1:1-1:10 HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3 | H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l),<br>HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G;<br>HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK;<br>H3FF; H3FJ   |
|   | Immunogen:  |
|   | Peptide sequence around site of Phospho-Ser (28) derived from Human Histone H3.1.   |
|   | Storage:  |

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



Immunofluorescence staining of Hela cells with PACO58653 at 1:5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Chromatin Immunoprecipitation Hela (4\*10<sup>6</sup>, treated with 100nM calyculin A for 60min) were treated with Micrococcal Nuclease, sonicated, and immunoprecipitated with 5µg anti-HIST1H3A (PACO58653) or a control normal rabbit IgG. The resulting ChIP DNA was quantified using real-time PCR with primers against the beta - Globin promoter.