
## **NOP53 Antibody**

## PACO59493



Product Information	
Size:	Protein Background:
50ug	
Reactivity:	Gene ID:
Human, Rat	NOP53
Source:	Uniprot
Rabbit	Q9NZM5
_	
lsotype:	Synonyms:
lsotype: lgG	Ribosome biogenesis protein NOP53 (Glioma tumor suppressor candidate region gene
lgG	Ribosome biogenesis protein NOP53 (Glioma tumor suppressor candidate region gene 2 protein) (Protein interacting with carboxyl terminus 1) (PICT-1) (p60), NOP53, GLT
lgG Applications:	Ribosome biogenesis protein NOP53 (Glioma tumor suppressor candidate region gene 2 protein) (Protein interacting with carboxyl terminus 1) (PICT-1) (p60), NOP53, GLT GLTSCR2 PICT1
IgG Applications: ELISA, WB, IHC	Ribosome biogenesis protein NOP53 (Glioma tumor suppressor candidate region gene 2 protein) (Protein interacting with carboxyl terminus 1) (PICT-1) (p60), NOP53, GLT GLTSCR2 PICT1 Immunogen:



Western Blot. Positive WB detected in: MCF-7 whole cell lysate, Rat heart tissue. All lanes: NOP53 antibody at  $4.8\mu$ g/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 55 kDa. Observed band size: 55 kDa.

IHC image of PACO59493 diluted at 1:600 and staining in paraffinembedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.